

PubMed



Display Settings: Abstract

FREE FINAL TEXT
OXFORD JOURNALS

Nephrol Dial Transplant. 2010 Aug;25(8):2730-7. doi: 10.1093/ndt/gfq082. Epub 2010 Feb 26.

Electrolysed-reduced water dialysate improves T-cell damage in end-stage renal disease patients with chronic haemodialysis.

Huang KC, Hsu SP, Yang CC, Ou-Yang P, Lee KT, Morisawa S, Otsubo K, Chien CT.

Department of Family Medicine, National Taiwan University College of Medicine and National Taiwan University Hospital, Taipei, Taiwan.

Abstract

BACKGROUND: T-cell damage by increased oxidative stress in end-stage renal disease (ESRD) patients undergoing chronic haemodialysis (HD) led to the increased T-cell apoptosis and the alteration of surface markers and Th1/Th2 ratio in CD4(+) T lymphocytes. Antioxidant electrolysed-reduced water (ERW) was used as the dialysate in ESRD patients undergoing chronic HD to test for improved oxidative stress-related T-cell apoptosis, alterations of surface markers and intracellular cytokine profile.

METHODS: We evaluated apoptosis formation by annexin V, CD25-related surface markers, and cytokine ratio of Th1/Th2 in CD4(+) T lymphocytes and Tc1/Tc2 in CD8(+) T lymphocytes of 42 ESRD patients haemodialysed with ERW for 1 year.

RESULTS: In comparison to 12 healthy individuals, the ESRD patients had more T-cell apoptosis and less CD3(+), CD4(+) and CD8(+) T cells and CD25/CD69/CD94/CD3(+) phenotypes at baseline. Lower intracellular IL-2 and IFN-gamma levels in the Th1/CD4(+) and Tc1/CD8(+) cells and higher intracellular IL-4, IL-6 and IL-10 levels in the Th2/CD4(+) and Tc2/CD8(+) cells were also noted in the ESRD patients. After a 1-year ERW treatment, the patients had a decrease in T-cell apoptosis and increases in CD3(+), CD4(+) and CD8(+) cell numbers and CD25/CD69/CD94/CD3(+) phenotypes in the T cells. The intracellular IL-2 and IFN-gamma levels in the Th1/Tc1 cells significantly ($P < 0.05$) increased and the intracellular IL-4, IL-6 and IL-10 levels in the Th2/Tc2 cells decreased. Furthermore, the Th1/Th2 and Tc1/Tc2 cytokine ratios were improved toward a normal status.

CONCLUSION: One-year ERW treatment effectively ameliorated T-cell apoptosis, altered CD25-related surface markers and intracellular cytokine profile in the HD patients.

PMID: 20190245 [PubMed - indexed for MEDLINE] [Free full text](#)

Publication Types, MeSH Terms, Substances

Publication Types

[Evaluation Studies](#)

[Research Support, Non-U.S. Gov't](#)

MeSH Terms

[Adult](#)

[Aged](#)[Aged, 80 and over](#)[Apoptosis*/drug effects](#)[C-Reactive Protein/metabolism](#)[CD4-Positive T-Lymphocytes/immunology](#)[CD4-Positive T-Lymphocytes/pathology*](#)[CD8-Positive T-Lymphocytes/immunology](#)[CD8-Positive T-Lymphocytes/pathology](#)[Case-Control Studies](#)[Dialysis Solutions/pharmacology](#)[Dialysis Solutions/therapeutic use*](#)[Female](#)[Humans](#)[Hydrogen Peroxide/pharmacology](#)[Hydrogen Peroxide/therapeutic use*](#)[Interleukin-2 Receptor alpha Subunit/metabolism](#)[Interleukin-6/metabolism](#)[Kidney Failure, Chronic/blood](#)[Kidney Failure, Chronic/pathology](#)[Kidney Failure, Chronic/therapy*](#)[Longitudinal Studies](#)[Male](#)[Middle Aged](#)[Renal Dialysis*](#)[Th1 Cells/pathology](#)[Th2 Cells/pathology](#)[Treatment Outcome](#)**Substances**[Dialysis Solutions](#)[Interleukin-2 Receptor alpha Subunit](#)[Interleukin-6](#)[Superoxidized water](#)[Hydrogen Peroxide](#)[C-Reactive Protein](#)**LinkOut - more resources****Full Text Sources**[HighWire](#)[EBSCO](#)[OhioLINK Electronic Journal Center](#)[Ovid Technologies, Inc.](#)**Medical**[Blood - MedlinePlus Health Information](#)

[Chronic Kidney Disease - MedlinePlus Health Information](#)

[Dialysis - MedlinePlus Health Information](#)

[Kidney Failure - MedlinePlus Health Information](#)

Molecular Biology Databases

[HYDROGEN PEROXIDE - HSDB](#)

Miscellaneous

[NCI CPTC Antibody Characterization Program](#)